

Hygeia: journal for drugs and medicines

April 2013 - September 2013

OPEN ACCESS

A half yearly scientific, international, open access journal for drugs and medicines

Research article section: Natural products/short communication

Researcher ID: K-6431-2012 Article ID- Hygeia.J.D.Med/83/13



Protective effect of DHC-1, a Polyherbal formulation, against CCl₄ induced Liver damage.

Bafna PA¹ and Balaraman R^{2*}

1. Rayat Institute of Pharmacy, Rayat and Bahra Campus, Railmajra, Dist. Nawanshahar, Punjab – 144 533 INDIA.

2. Department of Pharmacy, Sumandeep University, A/P Pipariya, (Waghodia), Vadodara-391760, Gujarat, INDIA.

Article history: Received: 15November 2012, revised: 21December 2012, accepted: 17January 2013, Available online: 3 April 2013

Abstract

Plan: DHC-1, a standardized polyherbal formulation, was investigated for its hepatoprotective and antioxidant effects against CCl₄-induced liver damage in rats.

Methodology: DHC-1 at various doses (125, 250, 500 and 1000 mg/kg, p.o.) was studied for its effect on the levels of SGPT, SGOT, alkaline phosphatase and total bilirubin. The drug was also evaluated for its effect on markers of tissue oxidative stress, namely extent of lipid peroxidation (MDA), levels of enzymatic antioxidants (SOD and catalase), non-enzymatic antioxidant (GSH) and membrane bound enzymes (Ca²⁺ATPase, Mg²⁺ATPase and Na⁺K⁺ATPase).

Outcome: Reduction in the levels of serum markers of liver damage; and decrease in tissue MDA levels and increase in SOD, catalase, GSH and membrane bound enzymes indicated the hepatoprotective and antioxidant property of DHC-1. Thus it is evident that DHC-1, at least partly by virtue of its antioxidant activity, elicited protective effects on the liver against the oxidative damage induced by CCl₄.

Keywords: Polyherbal formulation; Hepatoprotective; CCl₄; lipid peroxidation, superoxide dismutase, catalase, reduced glutathione.

1. Introduction

Liver is an organ of paramount importance as it plays an essential role in maintaining the biological equilibrium of vertebrates. It is the "alchemical wizard" of the body, transforming toxins into harmless chemicals for excretion, and digestively absorbed nutrients into the proper biochemical forms which the cells can use to function. Yet the liver is probably the organ most assaulted by toxic modern lifestyles, full of pollution, stress, junk foods, drugs, etc. About five percent of all deaths worldwide are the result of liver diseases. It ranks ninth in overall causes of death in the U.S.¹ and is the fifth 'big killer' in England & Wales.



For Correspondence: Email: rbalaraman2000@yahoo.com, Contact: 09898183938

Hygeia.J.D.Med. Vol.5 (1), April 2013

© 2013, Hygeia journal for drugs and medicines, All rights reserved. 2229 3590, 09756221

Researcher ID: K-6431-2012

Treatment options for common liver diseases such as cirrhosis, fatty liver and chronic hepatitis are problematic. The effectiveness of treatments such as interferon, colchicine, penicillamine and corticosteroids are inconsistent at best and the incidence of side effects is profound. All too often the treatment is worse than the disease.² Physicians and patients are thus in need of effective therapeutic agents with a low incidence of side effects. Plants potentially constitute such a group; and thus there is a worldwide trend to go back to traditional medicinal plants. Many natural products of herbal origin are in use for the treatment of liver ailments.³⁻⁵

The pathogenesis of hepatic diseases as well as the role of oxidative stress and inflammation therein is well recognized,⁶ and consequently, blocking or retarding the chain reactions of oxidation and inflammation development could be promising therapeutic strategies for the prevention and treatment of liver injury. Search for crude drugs of plant origin with antioxidant activity has thus become a central focus for study of hepatoprotection today.

The hepatoprotective effects of *Bacopa monnieri*,⁷ *Emblia officinalis*,^{8,9} *Glycyrrhiza glabra*,^{10,11} *Mangifera indica*¹² and *Syzygium aromaticum*^{13,14} have been mentioned. Also, the antioxidant properties of these plants, namely *Bacopa monnieri*,^{15,16} *Emblia officinalis*,¹⁷⁻¹⁹ *Glycyrrhiza glabra*,^{20,21} *Mangifera indica*,²²⁻²⁴ and *Syzygium aromaticum*^{25,26} have been investigated earlier and were found to possess free radical scavenging properties.

The antioxidant activity²⁷ and tissue protective effects of DHC-1, a formulation containing the above mentioned plants, have already been proved in different oxidative stress-induced disease conditions like pylorus-ligation and ethanol-induced ulcers,²⁸ cisplatin-induced nephrotoxicity and isoproterenol-induced myocardial infarction²⁹ in rats.

The present study was thus aimed to further test the efficacy of DHC-1 against hepatic injury induced by CCl₄ in rats to determine the possible use of this formulation in preventing hepatic damage and to justify whether the formulation exerts hepatoprotective effects by means of its antioxidant activity.

2. Materials and Methods

2.1 Plant Material

Bacopa monnieri, *Emblia officinalis*, *Glycyrrhiza glabra*, *Mangifera indica* and *Syzygium aromaticum* were procured from a local supplier and identified by Dr. Kannan, Ph.D., Botanist, The Himalaya Drug Company, Bangalore. Samples were retained for reference purpose at the R & D herbarium. The HPTLC fingerprint analysis of the individual ingredients and the formulation (DHC-1) have already been reported²⁷.

2.2. Composition

Each gram of DHC-1 contains extracts of:

<i>Herbs</i>	<i>Voucher code</i>	<i>Part used</i>	<i>Qty.</i>
<i>Bacopa monnieri</i> Linn. (Scrophulariaceae)	HDHB-157	Whole plant	200 mg
<i>Emblica officinalis</i> Gaertn. (Euphorbiaceae)	HDHB-143	Fruit	200 mg
<i>Glycyrrhiza glabra</i> Linn. (Papilionaceae)	HDHB-174	Roots	200 mg
<i>Mangifera indica</i> Linn. (Anacardiaceae)	HDHB-17	Bark	200 mg
<i>Syzygium aromaticum</i> Linn. (Myrtaceae)	HDHB-208	Flower bud	200 mg

2.3. Animals:

Albino rats of Wistar strain weighing 150-180g were housed in polypropylene cages under standard light/dark cycle, with food and water provided *ad libitum*. The experimental protocol was approved by the Institutional Animal Ethical Committee (IAEC) and conducted according to the guidelines of the Committee for the Purpose of Control and Supervision of Experiments on Animals (CPCSEA), Animal Welfare Division, Ministry of Forests and Environment, Govt. of India, New Delhi (India).

2.4. Experimental Procedure:

The animals were divided into six groups each consisting of six rats. Group 1 represented the Normal control group, which received 5 ml/kg of vehicle (1% gum acacia) orally for a period of 15 days. Group 2 served as the Negative control and received the vehicle orally (1% gum acacia; 5ml/kg) for 15 days followed by administration of CCl₄ (2.5 ml/kg, p.o.) in olive oil (1:1). Groups 3-6 received DHC-1 orally at the doses of 125, 250, 500 and 1000 mg/kg, respectively for 15 days followed by CCl₄ administration. After 24 hours of CCl₄ administration, blood was collected and serum was separated for estimations of SGPT, SGOT, alkaline phosphatase and total bilirubin using standard diagnostic kits [Span Diagnostics Ltd., Surat, India]. The animals were then sacrificed and the liver was dissected out, weighed and homogenized in chilled Tris buffer (10 mM, pH 7.4) at a concentration of 10% (w/v). The homogenates were then centrifuged at 10,000 x g at 0°C for 20 min using Remi C-24 high speed cooling centrifuge. The clear supernatant was then used for the assays of lipid peroxidation (MDA content), endogenous antioxidant enzymes like superoxide dismutase (SOD) and catalase (CAT), and reduced glutathione (GSH). The sediment was resuspended in ice-cold Tris buffer (10 mM, pH 7.4) to get a final concentration of 10% and was used for the estimation of different membrane bound enzymes (Na⁺K⁺ATPase, Ca²⁺ATPase, and Mg²⁺ATPase) and proteins.

2.5. Biochemical estimations:

Superoxide dismutase was determined by the method of Mishra and Fridovich.³⁰ Catalase was estimated by the method of Hugo Aebi as given by Colowick et al.³¹ Reduced glutathione was determined by the method of Moron et al.³² Lipid peroxidation or malondialdehyde formation was estimated by the method of Slater and Sawyer.³³

Membrane bound enzymes namely Na⁺K⁺ATPase, Ca²⁺ATPase and Mg²⁺ATPase were assayed according to the methods of Bonting,³⁴ Hjerken and Pan,³⁵ and Ohinishi et al.,³⁶ respectively. The inorganic phosphorus was estimated by the method of Fiske and Subarow.³⁷ Total proteins were determined by the method of Lowry et al.³⁸

2.6. Statistical Analysis of Data:

Results of all the above estimations have been indicated in terms of Mean \pm SEM. Difference between the groups was statistically determined by analysis of variance (ANOVA) followed by Tukey-Kramer Multiple Comparisons test, with the level of significance set at $p < 0.05$.

Table 1: Effect of DHC-1 on the serum levels of SGPT, SGOT, alkaline phosphatase and total bilirubin in CCl₄-induced hepatotoxicity in rats.

Groups	SGPT (U/ml)	SGOT (U/ml)	AlkP (mg/dl)	T.Bil. (mg/dl)
Group 1	47.88 \pm 2.33	77.64 \pm 2.85	62.50 \pm 2.57	0.103 \pm 0.009
Group 2	257.90 \pm 4.85 ^{***}	314.33 \pm 6.66 ^{***}	248.12 \pm 23.29 ^{***}	1.420 \pm 0.208 ^{***}
Group 3	224.12 \pm 8.83 ^{NS}	222.06 \pm 21.87 ^{NS}	163.37 \pm 7.55 ^{**}	0.807 \pm 0.099 ^{**}
Group 4	194.66 \pm 23.49 ^{NS}	221.47 \pm 41.43 ^{NS}	141.89 \pm 5.66 ^{**}	0.350 \pm 0.068 ^{***}
Group 5	166.94 \pm 22.66 [*]	209.26 \pm 11.15 [*]	139.51 \pm 10.61 ^{***}	0.143 \pm 0.009 ^{***}
Group 6	105.32 \pm 21.68 ^{***}	201.43 \pm 20.71 [*]	139.07 \pm 17.09 ^{***}	0.110 \pm 0.010 ^{***}
F value	22.127	12.310	20.429	28.942
P value	<0.0001	0.0002	<0.0001	<0.0001

Values are expressed as Mean \pm SEM. , Group 2 was compared with Group 1..Groups 3, 4, 5 and 6 were compared with Group 2.

* $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$; NS = Non Significant

Table 2: Effect of DHC-1 on the levels of lipid peroxidation (MDA content), reduced glutathione, superoxide dismutase and catalase in liver of rats in CCl₄-induced hepatotoxicity model.

Groups	Lipid Peroxidation (nmoles of MDA/mg protein)	Reduced Glutathione (μ g of GSH/mg protein)	Superoxide Dismutase (Units/mg protein)	Catalase (μ moles of H ₂ O ₂ consumed/min/mg protein)
Group 1	1.530 \pm 0.052	3.760 \pm 0.095	3.103 \pm 0.085	3.757 \pm 0.200
Group 2	5.331 \pm 0.102 ^{***}	0.474 \pm 0.035 ^{***}	1.497 \pm 0.084 ^{***}	1.520 \pm 0.161 ^{***}
Group 3	4.336 \pm 0.496 ^{NS}	1.854 \pm 0.177 ^{NS}	1.833 \pm 0.074 ^{NS}	1.630 \pm 0.129 ^{NS}
Group 4	4.029 \pm 0.154 [*]	3.074 \pm 0.590 ^{**}	2.050 \pm 0.110 [*]	1.883 \pm 0.074 ^{NS}
Group 5	2.555 \pm 0.166 ^{***}	4.291 \pm 0.386 ^{***}	2.177 \pm 0.077 ^{**}	2.260 \pm 0.055 [*]
Group 6	1.587 \pm 0.311 ^{***}	4.474 \pm 0.397 ^{***}	2.507 \pm 0.112 ^{***}	2.627 \pm 0.057 ^{***}
F value	36.297	20.897	37.224	43.980
P value	<0.0001	<0.0001	<0.0001	<0.0001

Values are expressed as Mean \pm SEM. , Group 2 was compared with Group 1. , Groups 3, 4, 5 and 6 were compared with Group 2.

* $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$; NS = Non Significant

Table 3: Effect of DHC-1 on the levels of Na⁺K⁺ATPase, Ca²⁺ATPase and Mg²⁺ATPase in liver of rats in CCL₄-induced hepatotoxicity model.

Groups	Na ⁺ K ⁺ ATPase (μ moles of inorganic phosphorus liberated/min/ mg protein)	Ca ²⁺ ATPase (μ moles of inorganic phosphorus liberated/min/ mg protein)	Mg ²⁺ ATPase (μ moles of inorganic phosphorus liberated/min/ mg protein)
Group 1	4.683 \pm 0.137	3.263 \pm 0.061	3.130 \pm 0.070
Group 2	2.500 \pm 0.085 ^{***}	1.555 \pm 0.039 ^{***}	2.166 \pm 0.082 ^{***}
Group 3	2.637 \pm 0.090 ^{NS}	1.704 \pm 0.098 ^{NS}	2.326 \pm 0.037 ^{NS}
Group 4	2.877 \pm 0.107 ^{NS}	2.136 \pm 0.091 [*]	2.458 \pm 0.120 ^{NS}
Group 5	3.270 \pm 0.201 ^{**}	2.477 \pm 0.065 ^{***}	2.577 \pm 0.108 ^{NS}
Group 6	3.977 \pm 0.067 ^{***}	3.019 \pm 0.170 ^{***}	2.837 \pm 0.081 ^{**}
F value	48.174	51.031	16.290
P value	<0.0001	<0.0001	<0.0001

Values are expressed as Mean \pm SEM. , Group 2 was compared with Group 1. ,Groups 3, 4, 5 and 6 were compared with Group 2.
*p<0.05; **p<0.01; ***p<0.001; NS = Non Significant

3. Results

Effect of DHC-1 on Serum Parameters:

Administration of CCl₄ in Group 2 animals (Negative control) resulted in a significant (p<0.001) elevation in SGPT, SGOT, alkaline phosphatase and total bilirubin levels, as compared to the Normal control group (Group 1).

The administration of DHC-1 significantly decreased the levels of SGPT and SGOT at the doses of 500 and 1000 mg/kg; whereas the alkaline phosphatase and total bilirubin were reduced at all the doses of DHC-1 (Table 1) as compared to the Negative control group (Group 2).

Effect of DHC-1 on Tissue Antioxidant Parameters:

In the liver of animals of Negative control group (Group 2), significant (p<0.001) reduction in the levels of SOD, catalase and reduced glutathione, and significant (p<0.001) increase in lipid peroxidation was observed when compared to Group 1 (Normal control) animals. Administration of DHC-1 at the doses of 250, 500 and 1000 mg/kg to animals of Groups 4, 5 and 6, respectively significantly increased the levels of SOD and GSH and decreased the levels of lipid peroxidation as compared to Group 2. A significant increase in the catalase level was observed at the doses of 500 mg/kg (Group 5) and 1000 mg/kg (Group 6) of DHC-1 as compared to Group 2 (Table 2).

In the negative control group animals (Group 2) significant ($p < 0.001$) decrease in the activities of membrane bound enzymes, namely $\text{Na}^+\text{K}^+\text{ATPase}$, $\text{Ca}^{2+}\text{ATPase}$ and $\text{Mg}^{2+}\text{ATPase}$ were observed as compared to the Normal control group (Group 1). The drug, DHC-1 was found to increase the activity of all the ATPases, namely $\text{Na}^+\text{K}^+\text{ATPase}$ (500 and 1000 mg/kg), $\text{Ca}^{2+}\text{ATPase}$ (250, 500 and 1000 mg/kg) and $\text{Mg}^{2+}\text{ATPase}$ activity (1000 mg/kg) at different doses (Table 3) as compared to the Negative control (Group 2).

4. Discussion

The presence of enzymes of the electron transport systems and high amounts of unsaturated fatty acids in the liver makes it vulnerable to peroxidative attack.³⁹

The hepatotoxin generally used to study the liver protective effect of drugs is CCl_4 because CCl_4 -induced liver dysfunction in rats simulates liver cirrhosis in man.^{40,41} It has been stated that one of the principal causes of carbon tetrachloride (CCl_4)-induced hepatopathy is lipid peroxidation by CCl_3^\bullet , a free radical derivative of the toxin.

Ko et al.⁴² have examined the impairment in hepatic antioxidant status during the development of CCl_4 -induced hepatotoxicity in rats and the protection of such tissue injury by pretreatment with vitamin E, herbal extracts or herbal preparations known to possess antioxidant activities. This generalized impairment in hepatic antioxidant defense mechanism was paralleled by an elevation in SGPT and SGOT activity, an indication of hepatocellular damage.⁴³

In liver injury due to hepatotoxin, there is a defective excretion of bile by the liver, which is reflected in the increased levels of alkaline phosphatase in serum.⁴⁴ Hyperbilirubinaemia is a very sensitive test to substantiate the functional integrity of the liver and severity of necrosis which increases the binding, conjugating and excretory capacity of hepatocytes that is proportional to the erythrocyte degeneration rate.⁴⁵

The stimulation of lipid peroxidation in either artificial membrane of liposomes or in subcellular organelles has been shown to increase membrane rigidity. In addition to the changes in fluidity, lipid peroxidation causes an increase in the ionic permeability and affects the surface potentials of the membranes. It has also been reported that administration of CCl_4 resulted in decrease in the activities of membrane bound enzymes⁴⁶⁻⁴⁸ ($\text{Na}^+\text{K}^+\text{ATPase}$, $\text{Ca}^{2+}\text{ATPase}$ and $\text{Mg}^{2+}\text{ATPase}$), thus leading to oxidative stress.

Administration of CCl_4 alone resulted in a significant elevation in SGPT, SGOT, alkaline phosphatase and total bilirubin levels. It also increased the levels of lipid peroxidation, and reduced the levels of both, endogenous antioxidants (SOD, catalase and GSH) and membrane bound enzymes ($\text{Na}^+\text{K}^+\text{ATPase}$, $\text{Ca}^{2+}\text{ATPase}$ and $\text{Mg}^{2+}\text{ATPase}$).

Depletion of elevated bilirubin level together with the suppression of the activities of SGPT, SGOT and ALP in the serum of rats treated with DHC-1, suggests the possibility of the herbal product to stabilize biliary dysfunction of rat liver during chronic injury with CCl₄. Reduction in lipid peroxidation (MDA) and enhancement in the levels of endogenous antioxidants (glutathione, catalase and SOD) proves the efficacy of DHC-1 in preventing free-radical induced damage in liver by CCl₄. Pretreatment with DHC-1 also increased the activity of all the ATPases indicating its membrane stabilizing action.

Thus the results prove that DHC-1 protected the liver from the damaging effects of CCl₄ by its antioxidant mechanism of action and can thus be used as a hepatoprotectant against such chemical insults.

The hepatoprotective effect of DHC-1 can be due to the ingredients, *Bacopa monnieri*,⁷ *Emblica officinalis*,^{8,9} *Glycyrrhiza glabra*,^{10,11} *Mangifera indica*¹² and *Syzygium aromaticum*^{13,14} which have individually been shown to protect the liver from the toxic effects of various hepatotoxicants.

As regards to the antioxidant activity of the constituents present in DHC-1, various reports are available. The *in vitro* antioxidant activity of *B. monniera*, one of the ingredients of DHC-1 was evaluated earlier by FeSO₄ induced lipid peroxidation in rat brain homogenate and the mechanism of action was thought to be through metal chelation at the initiation level and also as a chain breaker.¹⁵ The active tannoid principles of *Emblica officinalis* (amla), another ingredient of DHC-1, were found to induce an increase in both frontal cortical and striatal concentrations of the oxidative free radical scavenging enzymes, SOD, catalase and glutathione peroxidase and concomitant decrease in lipid peroxidation in these brain areas.¹⁷ Wang and Han¹⁰ suggested that the anti-lipid peroxidation effect of Glycyrrhiza flavonoids contributed to its protective action against carbon tetrachloride-induced hepatotoxicity. The extract of *Mangifera indica* reduces ischemia-induced neuronal loss and oxidative damage in the gerbil brain most probably due to the antioxidant activity of the extract.²³ The antioxidant activity of the extract was also studied on hydroxyl-mediated oxidation of bovine serum albumin (BSA) and in a hepatic microsome system and was found to reduce the oxidation of BSA and inhibited lipid peroxidation, which was, initiated enzymatically by NADPH.

The results suggested that the extract has an antioxidant activity probably due to its ability to scavenge free radicals involved in microsome lipid peroxidation. In addition, the extract's antioxidant profile *in vitro* was probably similar to its principal polyphenolic component, mangiferin, a glycosylated xanthone.²⁴

In another study²², Vimang, an aqueous extract of *M.indica* was found to provide significant protection against 12-O-tetra-decanoyl-phorbol-13-acetate (TPA)-induced oxidative damage in serum, liver, brain as well as against hyperproduction of ROS by peritoneal macrophages.

Thus Vimang could be useful to prevent the production of ROS and the oxidative tissue damages *in vivo*. The aroma extracts and aroma components isolated from *Syzygium aromaticum* (clove) were found to inhibit malondialdehyde formation from blood plasma oxidised with Fenton's reagent.⁴⁹

5. Conclusion

The results obtained from this study indicate that DHC-1 pretreatment offers significant protection to liver (hepatoprotective effect) and reduces the risk of CCl₄-induced liver damage by inhibiting lipid peroxidation and activating antioxidant defense system in the organ. The formulation may thus occupy a promising place in the market as a hepatoprotectant along with other popular herbal preparations like Liv 52, Tefroliv and Stimuliv, providing clinical studies also prove the same.

Earlier studies have proved the tissue protective effects of DHC-1 in other organs like heart, kidneys and gastric tissue and thus can be marketed as an overall good antioxidant formulation for defense against oxidative stress induced disease conditions.

Acknowledgement:

The authors are thankful to The Head, Pharmacy Dept., The M.S. University of Baroda, Vadodara (Gujarat) for providing all the facilities required to carry out this research work. The authors also extend their thanks to Himalaya Drug Co., Bangalore for providing the formulation, DHC-1; and UGC for providing financial assistance for successfully carrying out this work.

References

1. Bell BP, Navarro VJ, Manos MM, Murphy RC, Leyden WA, St. Louis TE, Kunze K, The epidemiology of newly diagnosed chronic liver diseases in the United States: findings of population-based sentinel surveillance. *Hepatology* 34, **2001**, 468A.
2. Scott Luper ND, A Review of Plants Used in the Treatment of Liver Disease: Part 1, *Altern Med Rev* 3, 6, **1998**, 410-421.
3. Venkateswaran S, Pari L, Viswanathan P, Menon VP, Protective effect of Livex, a herbal formulation against erythromycin estolate-induced hepatotoxicity in rats, *J Ethnopharmacol* 57, **1997**, 161-167.
4. Latha U, Rajesh MG, Latha MS, Hepatoprotective effect of an ayurvedic medicine, *Indian Drugs* 36, **1999**, 470-473.
5. Dhuley JN, Naik SR, Protective effect of Rhinax, a herbal formulation against CCl₄-induced liver injury and survival in rats, *J Ethnopharmacol* 56, **1997**, 159-164.
6. Tacke F, Luedde T, Trautwein C, Inflammatory pathways in liver homeostasis and liver injury, *Clin Rev Allerg Immu* 36, 1, **2009**; 4-12.
7. Rajalakshmy Menon B, Rathi MA, Thirumoorthi L, Gopalakrishnan VK, Potential effect of *Bacopa monnieri* on Nitrobenzene induced liver damage in rats, *Indian J Clin Biochem* 25, 4, **2010**, 401-404.
8. Jose JK, Kuttan R, Hepatoprotective activity of *Emblca officinalis* and Chyavanaprash, *J Ethnopharmacol* 72, 1-2, **2000**, 135-140.
9. Tasduq SA, Kaisar P, Gupta DK, Kapahi BK, Jyotsna S, Maheshwari HS, Johri RK, Protective effect of a 50% hydroalcoholic fruit extract of *Emblca officinalis* against anti-tuberculosis drugs induced liver toxicity, *Phytother Res* 19, 3, **2005**, 193-197.
10. Wang GS, Han ZW, The protective action of glycyrrhiza flavonoids against carbon tetrachloride hepatotoxicity in mice, *Yao Xue Xue Bao* 28, 8, **1993**, 572-576.
11. Yin G, Cao L, Xu P, Jeney G, Nakao M, Lu C, Hepatoprotective and antioxidant effects of *Glycyrrhiza glabra* extract against carbon tetrachloride (CCl₄)-induced hepatocyte damage in common carp (*Cyprinus carpio*), *Fish Physiol Biochem* 37, 1, **2011**, 209-216.
12. Pourahmad J, Eskandari MR, Shakibaei R, Kamalinejad M, A search for hepatoprotective activity of fruit extract of *Mangifera indica* L. against oxidative stress cytotoxicity, *Plant Foods Hum Nutr* 65, 1, **2010**, 83-89.
13. Rahman MKA, El-Megeid AAA, Hepatoprotective effect of soapworts (*Saponaria officinalis*), pomegranate peel (*Punica granatum* L) and cloves (*Syzygium aromaticum* linn) on mice with CCl₄ hepatic intoxication, *World Journal of Chemistry* 1, 1, **2006**, 41-46.
14. Prasad R, Ali S, Khan LA, Hepatoprotective effect of *Syzygium aromaticum* extract on acute liver injury induced by thioacetamide, *Int J Pharm Clin Res* 2, **2010**, 68-71.
15. Tripathi YB, Chaurasia S, Tripathi E, Upadhyay A, Dubey GP, *Bacopa monniera* Linn. as an antioxidant: mechanism of action, *Ind J Exp Biol* 34, **1996**, 520-526.

16. Ghosh T, Kumar Maity T, Das M, Bose A, Kumar Dash D, *In vitro* antioxidant and hepatoprotective activity of ethanolic extract of *Bacopa monnieri* Linn. aerial parts, *Iranian J Pharmacol Ther* 6, **2007**, 77-85.
17. Bhattacharya A, Chatterjee A, Ghosal S, Bhattacharya SK, Antioxidant activity of active tannoid principles of *E. officinalis*, *Ind J Exp Biol* 37, **1999**, 676-680.
18. Shukla V, Vashistha M, Singh S, Evaluation of antioxidant profile and activity of amalaki (*Embllica officinalis*), spirulina and wheat grass, *Ind J Clin Biochem* 24, 1, **2009**, 70-75.
19. Hazra B, Sarkar R, Biswas S, Manda N, Comparative study of the antioxidant and reactive oxygen species scavenging properties in the extracts of the fruits of *Terminalia chebula*, *Terminalia belerica* and *Embllica officinalis*, *BMC Complem Altern M* 10, **2010**, 20-35.
20. Hatano T, Fukuda T, Liu YZ, Noro T, Okuda T, Phenolic constituents of licorice. Correlation of phenolic constituents and licorice specimens from various sources and inhibitory effects of licorice extracts on xanthine oxidase and monoamine oxidase, *Yakugaku Zasshi* 111, **1991**, 311-321.
21. Rajesh MG, Latha MS, Protective activity of *Glycyrrhiza glabra* Linn. on carbon tetrachloride-induced peroxidative damage, *Ind J Pharmacol* 36, 5, **2004**, 284-287.
22. Sanchez GM, Re L, Giuliani A, Nunez-Selles AJ, Davison GP, Leon-Fernandez OS, Protective effects of *Mangifera indica* L. extract, mangiferin and selected antioxidants against TPA-induced biomolecules oxidation and peritoneal macrophage activation in mice, *Pharmacol Res* 42, **2000**, 565-573.
23. Martinez Sanchez G, Candelario-Jalil E, Giuliani A, Leon OS, Sam S, Delgado R, Nunez Selles AJ, *Mangifera indica* L. extract (QF808) reduced ischemia-induced neuronal loss and oxidative damage in the gerbil brain, *Free Radic Res* 35, 5, **2001**, 465-473.
24. Martinez G, Giuliani A, Leon OS, Perez G, Nunez Selles AJ, Effect of *Mangifera indica* L. extract (QF808) on protein and hepatic microsomal peroxidation, *Phytother Res* 15, 7, **2001**, 581-585.
25. Deans SG, Nobel RC, Hiltunen R, Wuryani W, Penzes LG, Antimicrobial and antioxidant properties of *Syzygium aromaticum*, *Flavour Fragr J* 10, **1995**, 323-327.
26. Makhuchit S, Antioxidant and Nitric Oxide Inhibition Activities of Thai Medicinal Plants, *J Med Assoc Thai* 93, Suppl.7, **2010**, S227-S235.
27. Bafna PA, Balaraman R, Antioxidant activity of DHC-1, a herbal formulation, *J Ethnopharmacol* 94, **2004a**, 135-141.
28. Bafna PA, Balaraman R, Anti-ulcer and antioxidant activity of DHC-1, an herbal formulation, *J Ethnopharmacol* 90, **2004b**, 123-127.
29. Bafna PA, Balaraman R, Antioxidant activity of DHC-1, an herbal formulation, in experimentally induced cardiac and renal damage, *Phytother Res* 19, 3, **2005**, 216-221.
30. Mishra HP, Fridovich I, The role of superoxide anion in the auto-oxidation of epinephrine and a simple assay for superoxide dismutase, *J Biol Chem* 247, **1972**, 3170-3175.
31. Colowick SP, Kaplan NO, Packer L, *Methods in Enzymology*, Academic Press, London, **1984**, p. 121-125.
32. Moron MS, Depierre JW, Mannervik B, Levels of glutathione, glutathione reductase and glutathione S-transferase activities in rat lung and liver, *Biochimica et Biophysica ACTA* 582, **1979**, 67-78.
33. Slater TF, Sawyer BC, The stimulatory effects of carbon tetrachloride and other halogenoalkanes on peroxidative reactions in rat liver fractions *in vitro*, *Biochem J* 123, **1971**, 805-814.
34. Bonting SL, *Presence of enzyme system in mammalian tissues. Membrane and Ion transport*, Wiley Inter Science, London, **1970**, p. 257-263.
35. Hjerken S, Pan H, Purification and characterization of two forms of a low affinity Ca²⁺-ATPase from erythrocyte membranes, *Biochimica et Biophysica ACTA* 728, **1983**, 281-288.
36. Ohinishi T, Suzuki T, Suzuki Y, Ozawa K, A comparative study of plasma membrane Mg²⁺-ATPase activities in normal, regenerating and malignant cells, *Biochimica et Biophysica ACTA* 684, **1982**, 67-74.
37. Fiske CH, Subbarow YT, Colorimetric determination of phosphorus, *J Biol Chem* 66, **1925**, 375-400.
38. Lowry OH, Rosenbrough NJ, Farr AC, Randell RJ, Protein measurement with folin-phenol reagent, *J Biol Chem* 193, **1955**, 265-275.
39. Poli G, Albano E, Dianzani MU, The role of lipid peroxidation in liver damage, *Chem Phys Lipids* 45, **1987**, 117-142.
40. Perez-Tamayo R, Is cirrhosis of the liver experimentally produced by CCl₄ an adequate model of human cirrhosis? *Hepatol* 3, **1983**, 112-120.
41. Wensing G, Sabra R, Branch RA, Renal and systemic hemodynamics in experimental cirrhosis in rats: relation to hepatic function, *Hepatol* 12, **1990**, 13-19.
42. Ko KM, Yick PK, Chiu TW, Hui TY, Cheng CHK, Kong YC, Impaired hepatic antioxidant status in carbon tetrachloride intoxicated rats: an *in vivo* model for screening herbal extracts with antioxidant activities, *Fitoterapia* LXIV, 6, **1993**, 539-544.
43. Naziroglu M, Cay M, Ustundag B, Aksakal M, Yekeller H, Protective effects of vitamin E on carbon tetrachloride-induced liver damage in rats, *Cell Biochem Funct* 17, **1999**, 253-259.
44. Rao RR, Mechanism of drug induced hepatotoxicity, *Ind J Pharmacol* 5, **1973**, 313-318.
45. Singh B, Saxena AK, Chandan BK, Anand KK, Suri OP, Suri KA, Satti NK, Hepatoprotective activity of verbenaol on experimental liver damage in rodents, *Fitoterapia* 69, **1998**, 135-140.
46. Long RM, Moore L, Elevated cytosolic calcium in rat hepatocytes exposed to carbon tetrachloride, *J Pharmacol Exp Ther* 238, **1986**, 186-191.
47. Katsumata T, Murata T, Yamaguchi M, Alteration in calcium content and Ca²⁺-ATPase activity in the liver nuclei of rats orally administered carbon tetrachloride, *Mol Cell Biochem* 185, 1-2, **1998**, 153-159.
48. Hemmings SJ, Pulga VB, Tran ST, Uwiera RR, Differential inhibitory effects of carbon tetrachloride on the hepatic plasma membrane, mitochondrial and endoplasmic reticular calcium transport systems: implications to hepatotoxicity, *Cell Biochem Funct* 20, 1, **2002**, 47-59.
49. Lee KG, Shibamoto T, Inhibition of malondialdehyde formation from blood plasma oxidation by aroma extracts and aroma components isolated from clove and eucalyptus, *Food Chem Toxicol* 39, 12, **2001**, 1199-1204.

Bafna PA, Balaraman R. Protective effect of DHC-1, a Polyherbal formulation, against CCL₄ induced Liver damage: *Hygeia.J.D.Med.* **2013**; 5(1):10-18. Available from <http://www.hygeiajournal.com/> / Article ID- Hygeia.J.D.Med/83/13